

## The Quaternary Range Dynamics of the Dwarf Lizard, *Parvilacerta parva* (Boulenger, 1887) (Squamata, Lacertidae) in the Anatolian Peninsula

Mehmet Kürşat Şahin<sup>1\*</sup>, Yusuf Kumlutaş<sup>2,3</sup>, Alexey Yanchukov<sup>4</sup>, Ortaç Çetintaş<sup>4</sup>, Kamil Candan<sup>2</sup>, Çetin Ilgaz<sup>2</sup>, Zafer Ayaş<sup>5</sup>

<sup>1</sup> Department of Biology, Kamil Özdağ Faculty of Science, Karamanoğlu Mehmetbey University, Karaman, Turkey

<sup>2</sup> Department of Biology, Faculty of Science, Dokuz Eylül University, İzmir, Buca, Turkey

<sup>3</sup> Research and Application Center for Fauna and Flora, Dokuz Eylül University, 35610, Buca-İzmir, Turkey

<sup>4</sup> Department of Biology, Faculty of Arts and Sciences, Bülent Ecevit University, Zonguldak, Turkey

<sup>5</sup> Department of Biology, Faculty of Science, Hacettepe University, Ankara, Turkey

\*Email: yasambilimci.kursat@gmail.com

Received: 08 September 2021 / Revised: 12 October 2021 / Accepted: 13 October 2021 / Published online: 14 October 2021. Ministry of Sciences, Research and Technology, Arak University, Iran.

**How to cite:** Şahin, M.K., Kumlutaş, Y., Yanchukov, A., Çetintaş, O., Candan, K., Ilgaz, Ç., Ayaş, Z (2021). The Quaternary Range Dynamics of the Dwarf Lizard, *Parvilacerta parva* (Boulenger, 1887) (Squamata, Lacertidae) in the Anatolian Peninsula, 6(X), XXX-XXX. <https://doi.org/10.22120/jwb.2021.540551.1259>

### Abstract

The dwarf lizard, *Parvilacerta parva*, is a characteristic member of the steppe biome in Irano - Anatolian biological hotspot. While this lizard has been included in local faunistic surveys and its morphological variation was addressed, no targeted study has been performed on the ecology and distribution of the species. Here we investigate the range dynamics of dwarf lizard during recent glacial and interglacial periods. We looked at the effects of climatic oscillations on species distribution at Present, Last Glacial Maximum (LGM), and Last Interglacial (LIG) periods using ecological niche modelling (ENM), based on our own fieldwork and literature data. The model

results suggest that the range of dwarf lizards contracted during the LIG and expanded during the LGM, opposite to the pattern observed in many other temperate reptiles. During the LIG, distribution of the dwarf lizards had been restricted to the mountainous steppe habitats in Northeastern Anatolia, but during the LGM it expanded to the west by including the new steppe habitats in Sultan, Emir and Murat mountains and adjacent areas. Climatic factors had a strong influence on shaping the spatiotemporal habitat. The Anatolian Biogeographic Region overlaps with Irano - Anatolian biodiversity hotspot, reflecting remarkable species richness in this area. However, faunal elements of the hotspots are under threat due to not only global climate change, but also anthropogenic pressures, such as habitat loss and overgrazing. Our results suggest that the dwarf lizards have a potential as indicators for tracking the local effects of global climate change as well as human induced degradation of the steppe habitat.

**Keywords:** last glacial maximum, last interglacial, ecological niche modelling, species distribution, Turkey

## **Introduction**

The real accessibility of geographically and climatologically supported organismal information proposes new perceptions into evolutionary processes (Rivera et al., 2021). One of the commonly used approaches to examine these viewpoints is to evaluate species distribution patterns via benefiting from ecological niche modelling (ENM) (Phillips et al., 2006). ENM is deeply associated with georeferenced occurrence data of species (set of geographical coordinates where the species is observed) and environmental data (i.e climatological, and/or topological data) and generates potential geographic distribution predictions for the species of interest (Guisan & Thuiller, 2005). Due to its continuously developing stages, this modelling approach not only gives the current potential distribution of the species, but also past and future distribution patterns by using reconstructed climate data projections (IPCC, 2007; Waltari et al., 2007).

One of the undeniable factors to understand the biogeography is that the modern-day populations and communities have been shaped by the interactions of organisms and their environments in the past (Gherghel, 2021). Besides, it is well known that the climatic oscillations in the Late Pleistocene glacial (Last Glacial Maximum: LGM 21000 years ago) and interglacial periods (the Last Interglacial: LIG ~130000-116000 years ago) have a profound influence on ecosystems in the Northern Hemisphere (Conradi et al., 2020; Perktaş, Gür, Sağlam, et al., 2015; Ülker et al., 2018).

The “expansion-contraction” model has been used widely and successfully to explain the change in species distribution during climate fluctuations (Horreo et al., 2018; Provan & Bennett, 2008). According to this model, the climate during the glacial periods was too cold and dry for the survival of most species within their present-day range, so they were retreating to warmer parts of the range at lower latitudes. Thereafter, the species recolonized the new areas at higher elevations with a range expansion during the interglacial periods. However, this model mostly applies to temperate species and is less suitable for those species that have preferences to drier and colder habitats (Gür, 2013).

Studies in the last two decades have uncovered the role of the Anatolian Peninsula as a refugium

for several taxa (Gür, 2013; Korkmaz et al., 2014; Kornilios et al., 2012; Perktaş et al., 2015a; Perktaş et al., 2015b; Rokas et al., 2003). Wetter conditions with a higher precipitation rate during the LGM along the western part of the peninsula (Sarıkaya et al., 2008) enabled wider suitable vegetation ranges for the species, which are adapted to the poikilothermic strategy. Multiple studies demonstrated that lizards are particularly good model organisms to examine the evolutionary history dynamics via ENM predictions in terrestrial habitats (Gül et al., 2018; Kurnaz et al., 2016). Because ENM has been widely used to assess geographic range of species via abiotic environmental conditions (i.e. temperature, precipitation dynamics, and topography) (Kozak et al., 2008). The reason behind this phenomenon was described as the fundamental niche of the species by (Hutchinson, 1957).

The Anatolian Biogeographic Region includes a steppe and steppe forest region of 321754 km<sup>2</sup>, however the actual range of the steppe vegetation is limited to 180316 km<sup>2</sup> (56% of the potential distribution). The rest of the region (44%) has been transformed to croplands (Ambarlı et al., 2016). The Anatolian steppes host 63% of the herpetofauna diversity in Turkey (Ambarlı et al., 2016). While the complex steppe biome includes a large number of species (Zeng et al., 2014), examining a single representative species can be a practical way for tracing the effects of climate on spatio-temporal distribution of the entire ecosystem.

The dwarf lizard, *Parvilacerta parva* (Boulenger, 1887), is a diurnally active species distributed along the steppes and alpine meadows of central and eastern Anatolia and adjacent Armenia. Its chorotype is described as Armeno-Anatolian endemic (Kurnaz, 2020; Sindaco et al., 2000). The geographic distribution of the dwarf lizard has two major climate zones, Central and North-Eastern Anatolia, that are drier and colder than the other zones in Turkey (Beck et al., 2018; Unal et al., 2003). Annual activity period of the dwarf lizards is between mid-April and the end of September, and they hibernate during the winter and very early spring (Arakelyan et al., 2011). This strong dependence on abiotic factors makes the dwarf lizards highly suitable organisms to study via ENM.

Our main aim is to evaluate the effects of climate oscillations on the distribution pattern of the dwarf lizard. We hypothesized that their distribution should have contracted during interglacial climatic conditions (LIG and present) and expanded during drier and colder conditions in LGM. Additionally, we speculate about the role of the Anatolian steppes in the distribution of dwarf lizards throughout the LIG and LGM.

## Material and methods

### Study area and input data

This study was conducted within 25-46° East Longitude and 34-43° North Latitude, covering the Anatolian Peninsula, the Caucasus Region (since part of the dwarf lizard range in Armenian steppe habitats) and the Levantine Region (where the sister species, *Parvilacerta fraasi*, occurs).

A total of 119 occurrence records were gathered from our field observations and the literature records (Atagün, 1984; Baran et al., 1992; Basoğlu & Baran, 1977; Kumlutaş et al., 2004; Mülayim et al., 2001; Peters, 1962) (Supplementary File 1). In cases where the locality information was not directly given in any GPS format, an online geographic system softwares (i.e. Google Earth Pro) were used to determine the most accurate location. The map of current distribution for the dwarf lizards is shown in Figure 1. In order to avoid the spatial sampling biases and misinterpretation of

the distribution map, these occurrence records were spatially rarefied with removing at least one locality in each 10 km by using SDM Toolbox 2.0 (Brown, 2014).



**Figure 1.** Species occurrence records for the dwarf lizard in Anatolia

Nineteen bioclimatic, and one topographic (elevation) parameters were downloaded from WorldClim as v.1.4 (Hijmans et al., 2005; [www.worldclim.org](http://www.worldclim.org)) in 2.5 Arc Second spatial resolution (~4.63 km at the equator) (Supplementary File 2). In addition to present climatological parameters, Model for Interdisciplinary Research on Climate (MIROC) was used for paleoclimatological data on LIG and LGM (Watanabe et al., 2010). The resolution of the LIG climatic data was 30'', therefore we converted these data to a resolution of 2.5' for their compatibility with Present and LGM.

All layers were clipped by using ArcGIS 10.6.1 (ESRI) to fit the study area. Correlation coefficients between variables were calculated in R 3.6.3 (R. Core Team, 2018) and highly correlated variable pairs ( $r \geq |0.8|$ ) were excluded from analysis for eliminating the adverse consequences from other parameters (Table 1).



## Ecological Niche Modelling (ENM)

We used maximum entropy machine learning algorithm (MaxEnt), v 3.4.1 (Phillips et al., 2017) to construct the models under Present conditions and projecting to LIG and LGM. MaxEnt is a useful and widely adopted tool that uses only presence records to get a high model performance even with low occurrence data (Merow et al., 2013). The following settings were applied to the candidate models: the convergence threshold is 0.00001, maximum number of iteration is 500, the regularization multipliers are 0.1, 0.2, 0.3,...1), five feature classes (hinge, threshold, product, quadratic, and linear). 20% of occurrence data and 8263 background points were used as test points. In order to check the importance of variables, the jackknife test was used and ten bootstrap replicates estimated statistical support for the constructed models (Elith et al., 2011). The power of the model was determined by the surface area under the receiver-operator (ROC) and area under curve (AUC) (Peterson et al., 2008; Phillips et al., 2017). The AUC is a score between 0.5 (no better than random prediction) and 1 (the flawless predicted model).

## Results

Pearson correlation coefficients among climatic and topographic layers and indicated that four parameters varied 8.5 % to 72.8 % can be taken into account for the species distribution model of dwarf lizards. These were Bio 2 (Mean Diurnal Range), Bio 8 (Mean Temperature of Wettest Quarter), Bio 9 (Mean Temperature of Driest Quarter) and Bio 15 (Precipitation Seasonality). The contributions of the variables are given in Table 2. Based on the results, Bio 9 made the largest contribution to species presence at 72.8 %, and Bio 15 made the lowest contribution at 8.5%. However, the possible effect of Bio 8 was ignored due to its irregularity pattern as a climatological parameter.

Estimated training AUC value (mean  $\pm$  SD) from rarefied 103 occurrence points was  $0.886 \pm 0.027$  indicating that AUC value is sufficiently high to keep in the analysis. Besides, the small SD for the mean AUC indicated that the ENM performance was sufficiently strong in response to variation in the occurrence records for training and test. Lastly, the model constructed from the test occurrence records performed better in each replicate compared to a random prediction ( $P < 0.001$ ).

**Table 2.** Contribution of variables to model prediction

Variable	Percent contribution	Permutation importance
Bio_9	72.8	79.6
Bio_2	17.8	15.2
Bio_15	8.5	4.1
Bio_8	0.9	1.2

The logistic outputs of the ENM for three time periods (LIG, LGM and the present) are shown on Figure 2 a-c. The model prediction for the present period was profoundly consistent with the actual distribution range of the dwarf lizard, also suggesting that the dwarf lizard is almost at equilibrium with the climate (Figure 2 a). This map includes the steppe vegetation and alpine meadows of the Anatolian Peninsula, especially in the central and eastern side, which the species preferably inhabit. However, even though the species range extended to western part of the Peninsula, the predicted range does not display a continuity pattern as the central and the eastern parts.




Figure 2. Habitat suitability predictions of the dwarf lizard for a. Present, b. LGM, c. LIG

## **Discussion**

Several biotic and abiotic factors have large effects on the species distribution (Peterson, 2011). Therefore, in order to perceive the responses of dwarf lizards to the climatic oscillations through the Late Quaternary glacial - interglacial cycles, we used ENM as a useful tool in analytical biology, to evaluate the range of species and develop hypotheses about their evolutionary history. One of the most useful properties of ENM is that it only requires species occurrences and environmental variables.

The Irano - Anatolian biodiversity hotspot is one of the three worldwide hotspots spanning the Anatolian peninsula (Mittermeier et al., 2011). The steppic parts of the Anatolian Biogeographic Region overlaps with this hotspot, in which hosts one of the seven identified grassland hotspots in the Palearctic Realm (Figure 3) (Dengler et al., 2014). Three factors contribute to high level of biodiversity in this region: (i) its biological intersection position among Asia, Europe and Africa continents (Kışlalıoğlu & Berkes, 1987), (ii) land-use history (Ambarlı & Bilgin, 2014) and (iii) wide amplitude of abiotic factors (geology, soil type, climate regimes etc.) (Şekercioğlu et al., 2011).


The reactions of dwarf lizards to the climate changes through the Late Quaternary epoch are comparable to that in other steppe-dependent species, such as Anatolian ground squirrels (Gür, 2013). The range of dwarf lizards contracted during the interglacial periods and expanded during the glacial periods and. In other words, the general pattern of inter-glacial expansion and glacial range contraction of the temperate species ranges (Hewitt, 1999; Hewitt, 2000) does not apply to dwarf lizards.

Although the phylogeography of the temperate and cold-adapted species have been comprehensively studied so far, the third group of species, like the dwarf lizard that are adapted to dry and occasionally warm conditions, and occupy steppes and other types of dry meadows, have received less attention (Kajtoch et al., 2016). Distributions of populations of mountainous and boreal taxa were narrower during the inter-glaciations than in glacials, including the Holocene epoch (Schmitt, 2009; Schönswetter et al., 2005; Varga, 2009). Importantly, steppe species are the most threatened organisms and steppes are the most threatened ecosystems in European continent due to natural (i.e. drying and regional rainfall shifts, enhancing fire risk and restraining the capacity of species to move), and to the human-induced degradation and fractionation of steppes (Cremene et al., 2005; Fekete et al., 2014; Janišová et al., 2011). Further studies in the overlapping areas of the Anatolian Biogeographic Region and the Irano-Anatolian hotspot might also point that these threats can be valid for these areas.

The fact that the predicted distribution of the dwarf lizards during the present period corresponds closely to its actual range, suggests that dwarf lizards are close to equilibrium with climatological dynamics (Figure 1, 2a). Moreover, this equilibrium also makes the reconstructed past predictions consistent, due to assumptions of species - climate equilibrium and stability of ecological niches in the course of time have been met (Nogués-Bravo, 2009). Bio-climatological envelope that was made by BIOs 2, 8, 9, 15 that are related to dryland ecosystems has met the expectations for a steppe habitat specialist species that includes the dwarf lizard. The Bio 9 refers to “mean temperature of driest quarter”, Bio 2 - “mean diurnal range”, and Bio 15 is “precipitation seasonality”. These three variables contribute ~ 99.1% to the distribution of the dwarf lizard, and

have also shaped the ecosystem services of habitats (Chen et al., 2015; Petrosyan et al., 2019). However, the possible effect of Bio 8 “Mean Temperature of Wettest Quarter” (percent contribution: 0.9%, permutation importance: 1.2 %) was ignored due to its odd spatial irregularities within the frame of discontinuities between adjoining pixels within the nonappearance of ecological gradients on the ground as a climatological parameter (Ashraf et al., 2017; Behroozian et al., 2020).

Anatolian steppes expanded remarkably as result of cold and dry conditions during the glacial periods (Atalay, 2020). It is most likely that dwarf lizards have followed this habitat expansion. For instance, the steppes in Sultan, Emir and Murat Mountains may have enabled the lizards to extend their distribution through the LGM (Figure 2b). However, the climatic conditions in the interglacial periods (LIG, present) caused contraction of dwarf lizard populations (Figure 2a, c).



**Figure 3.** The general land cover types of Turkey (modified from Corine 2018)

## Conclusion

Incidentally, as these lizards are currently in the contraction phase, their habitat is further deteriorating due to anthropogenic degradation, such as overgrazing (Fırıncioğlu et al., 2009). We argue that the dwarf lizard can be considered an indicator species for Anatolian dry steppe habitat. Further studies on the molecular biogeography of dwarf lizards will help to reveal more details of the history of range dynamics in this lizard species.

## Conflict of Interest

The authors declare that they have no conflict of interest.

## Acknowledgments

This study was based on Mehmet Kürşat Şahin’s PhD thesis supervised by Dr. Zafer Ayaş. We give

special thanks to Dr. David Tarkhnishvili for his comments and Dr. Murat Afsar, Dr. Mert Karış, Dr. Bahadır Akman, Doğan Sözbilen, Dr. Murat Tosunoğlu, Dr. Uğur Cengiz Erişmiş for sharing information on locality data. Finally, we wish to thank anonymous reviewers who helped us to improve the first draft. This work was supported by the Hacettepe University Scientific Research Coordination Unit. (Project Number FHD-2018-16241).

## References

- Ambarlı, D., & Bilgin, C. C. (2014). Effects of landscape, land use and vegetation on bird community composition and diversity in Inner Anatolian steppes. *Agriculture, Ecosystems & Environment*, 182, 37–46.
- Ambarlı, D., Zeydanlı, U. S., Balkız, Ö., Aslan, S., Karaçetin, E., Sözen, M., Ilgaz, Ç., Ergen, A. G., Lise, Y., & Çağlayan, S. D. (2016). An overview of biodiversity and conservation status of steppes of the Anatolian Biogeographical Region. *Biodiversity and Conservation*, 25(12), 2491–2519.
- Arakelyan, M. S., Danielyan, F. D., Corti, C., Sindaco, R., & Leviton, A. E. (2011). “Herpetofauna of Armenia and Nagorno-Karabakh. Society for the Study of Amphibians and Reptiles (SSAR).
- Ashraf, U., Peterson, A. T., Chaudhry, M. N., Ashraf, I., Saqib, Z., Rashid Ahmad, S., & Ali, H. (2017). Ecological niche model comparison under different climate scenarios: A case study of *Olea* spp. In Asia. *Ecosphere*, 8(5), e01825.
- Atagün, F. (1984). Türkiye’de *Lacerta parva* (Reptilia, Lacertidae)’nın Taksonomik Araştırılması. Ege Üniversitesi (in Turkish).
- Atalay, İ. (2020). Paleoenvironmental conditions of the Late Pleistocene and early Holocene in Anatolia, Turkey. In *Quaternary deserts and climatic change* (pp. 227–237). CRC Press.
- Baran, İ., Yılmaz, I., Kete, R., Kumlutas, Y., & Durmus, S. H. (1992). Orta Karadeniz Bölgesinin Herpetofaunasi. *Turkish Journal of Zoology*, 16(3), 275–288.
- Basoğlu, M., & Baran, I. (1977). Türkiye Sürüngenleri, Kısım I, Kaplumbağa ve Kertenkeleler [Turkish Reptiles. Part I. Turtles and Lizards]. Ege Üniversitesi Kitaplar Serisi, 76, 1–219 (in Turkish).
- Beck, H. E., Zimmermann, N. E., McVicar, T. R., Vergopolan, N., Berg, A., & Wood, E. F. (2018). Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Scientific Data*, 5, 180214.
- Behroozian, M., Ejtehadi, H., Peterson, A. T., Memariani, F., & Mesdaghi, M. (2020). Climate change influences on the potential distribution of *Dianthus polylepis* Bien. Ex Boiss.(Caryophyllaceae), an endemic species in the Irano-Turanian region. *PloS One*, 15(8), e0237527.
- Brown, J. L. (2014). SDM toolbox: A python-based GIS toolkit for landscape genetic, biogeographic and species distribution model analyses. *Methods in Ecology and Evolution*, 5(7), 694–700.

- Chen, L., Li, H. E., Zhang, P., Zhao, X., Zhou, L., Liu, T., Hu, H., Bai, Y., Shen, H., & Fang, J. (2015). Climate and native grassland vegetation as drivers of the community structures of shrub-encroached grasslands in Inner Mongolia, China. *Landscape Ecology*, 30(9), 1627–1641.
- Conradi, T., Van Meerbeek, K., Ordonez, A., & Svenning, J.-C. (2020). Biogeographic historical legacies in the net primary productivity of Northern Hemisphere forests. *Ecology Letters*, 23(5), 800–810.
- Corine Land Cover (2018), Available online: <https://land.copernicus.eu/pan-european/corine-land-cover/clc2018> Retrieved August 20, 2020
- Cremene, C., Groza, G., Rakosy, L., Schileyko, A. A., Baur, A., Erhardt, A., & Baur, B. (2005). Alterations of steppe-like grasslands in Eastern Europe: A threat to regional biodiversity hotspots. *Conservation Biology*, 19(5), 1606–1618.
- Dengler, J., Janišová, M., Török, P., & Wellstein, C. (2014). Biodiversity of Palaeartic grasslands: A synthesis. *Agriculture, Ecosystems & Environment*, 182, 1–14.
- Elith, J., Phillips, S. J., Hastie, T., Dudík, M., Chee, Y. E., & Yates, C. J. (2011). A statistical explanation of MaxEnt for ecologists. *Diversity and Distributions*, 17(1), 43–57.
- Fekete, G., Molnár, Z., Magyari, E., Somodi, I., & Varga, Z. (2014). A new framework for understanding Pannonian vegetation patterns: Regularities, deviations and uniqueness. *Community Ecology*, 15(1), 12–26.
- Fıncıoğlu, H. K., Seefeldt, S. S., Şahin, B., & Vural, M. (2009). Assessment of grazing effect on sheep fescue (*Festuca valesiaca*) dominated steppe rangelands, in the semi-arid Central Anatolian region of Turkey. *Journal of Arid Environments*, 73(12), 1149–1157.
- Gherghel, I. (2021). How Ecology and Evolution Shape Species Distributions and Ecological Interactions Across Time and Space. Case Western Reserve University.
- Guisan, A., & Thuiller, W. (2005). Predicting species distribution: Offering more than simple habitat models. *Ecology Letters*, 8(9), 993–1009.
- Gül, S., Kumlutaş, Y., & Ilgaz, Ç. (2018). Potential distribution under different climatic scenarios of climate change of the vulnerable Caucasian salamander (*Mertensiella caucasica*): A case study of the Caucasus Hotspot. *Biologia*, 73(2), 175–184.
- Gür, H. (2013). The effects of the Late Quaternary glacial–interglacial cycles on Anatolian ground squirrels: Range expansion during the glacial periods? *Biological Journal of the Linnean Society*, 109(1), 19–32.
- Hewitt, G. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, 405(6789), 907.
- Hewitt, G. (1999). Post-glacial re-colonization of European biota. *Biological Journal of the Linnean Society*, 68(1–2), 87–112.
- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 25(15), 1965–1978.

- Horreo, J. L., Pelaez, M. L., Suárez, T., Breedveld, M. C., Heulin, B., Surget-Groba, Y., Oksanen, T. A., & Fitze, P. S. (2018). Phylogeography, evolutionary history and effects of glaciations in a species (*Zootoca vivipara*) inhabiting multiple biogeographic regions. *Journal of Biogeography*, 45(7), 1616–1627.
- Hutchinson, G. E. (1957). Concluding remarks. *Cold Spring Harbor Symposium on Quantitative Biology*, 22, 415–457.
- IPCC. (2007). *Climate Change 2007: The Physical Science Basis: Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change: Summary for Policymakers and Technical Summary and Frequently Asked Questions*. Cambridge University Press.
- Janišová, M., Bartha, S., Kiehl, K., & Dengler, J. (2011). Advances in the conservation of dry grasslands: Introduction to contributions from the seventh European Dry Grassland Meeting. *Plant Biosystems-An International Journal Dealing with All Aspects of Plant Biology*, 145(3), 507–513.
- Kajtoch, Ł., Cieślak, E., Varga, Z., Paul, W., Mazur, M. A., Sramkó, G., & Kubisz, D. (2016). Phylogeographic patterns of steppe species in Eastern Central Europe: A review and the implications for conservation. *Biodiversity and Conservation*, 25(12), 2309–2339.
- Kışlalıoğlu, M., Berkes, F., (1987). *Biyolojik Çeşitlilik. Türkiye Çevre Sorunları Vakfı*, Ankara (in Turkish).
- Korkmaz, E. M., Lunt, D. H., Çıplak, B., Değerli, N., & Başıbüyük, H. H. (2014). The contribution of Anatolia to European phylogeography: The centre of origin of the meadow grasshopper, *Chorthippus parallelus*. *Journal of Biogeography*, 41(9), 1793–1805.
- Kornilios, P., Ilgaz, Ç., Kumlutaş, Y., Lymberakis, P., Moravec, J., Sindaco, R., Rastegar-Pouyani, N., Afroosheh, M., Giokas, S., & Fraguadakis-Tsolis, S. (2012). Neogene climatic oscillations shape the biogeography and evolutionary history of the Eurasian blindsnake. *Molecular Phylogenetics and Evolution*, 62(3), 856–873.
- Kozak, K. H., Graham, C. H., & Wiens, J. J. (2008). Integrating GIS-based environmental data into evolutionary biology. *Trends in Ecology & Evolution*, 23(3), 141–148.
- Kumlutaş, Y., Durmuş, S. H., Kaska, Y., Öz, M., & Tunç, M. R. (2004). A morphological and taxonomic study on *Lacerta parva* Boulenger, 1887 (Sauria: Lacertidae) from West Taurus, Turkey. *Asiatic Herpetological Research*, 10, 202–207.
- Kurnaz, M. (2020). Species list of Amphibians and Reptiles from Turkey. *Journal of Animal Diversity*, 2(4), 10–32.
- Kurnaz, M., Gül, S., Bülbül, U., & Kutrup, B. (2016). The potential distribution of *Darevskia derjugini* (Nikolsky, 1898) with new locality records from Turkey. *Developments in Science and Engineering*, St. Kliment Ohridski Univ. Press, Sofia, 189–195.
- Merow, C., Smith, M. J., & Silander Jr, J. A. (2013). A practical guide to MaxEnt for modeling species' distributions: What it does, and why inputs and settings matter. *Ecography*, 36(10), 1058–1069.

- Mittermeier, R. A., Turner, W. R., Larsen, F. W., Brooks, T. M., & Gascon, C. (2011). Global biodiversity conservation: The critical role of hotspots. In *Biodiversity hotspots* (pp. 3–22). Springer.
- Mülayim, A., Tok, C. V., & Ayaz, D. (2001). Beyşehir (Konya) Civarından Toplanan *Lacerta parva* Boulenger, 1887 (Sauria: Lacertidae) Örnekleri Üzerinde Morfolojik Bir Çalışma, *Anadolu University Journal of Science and Technology*, 2(2), 345-349 .
- Nogués-Bravo, D. (2009). Predicting the past distribution of species climatic niches. *Global Ecology and Biogeography*, 18(5), 521–531.
- Perktaş, U., Gür, H., & Ada, E. (2015a). Historical demography of the Eurasian green woodpecker: Integrating phylogeography and ecological niche modelling to test glacial refugia hypothesis. *Folia Zoologica*, 64(3), 284–295.
- Perktaş, U., Gür, H., Sağlam, İ. K., & Quintero, E. (2015b). Climate-driven range shifts and demographic events over the history of Kruper’s nuthatch *Sitta krueperi*. *Bird Study*, 62(1), 14–28.
- Peters, G. (1962). Die Zwergeidechse (*Lacerta parva*, Boulenger) und ihre Verwandtschaftsbeziehungen zu anderen Lacertiden, insbesondere zur Libanon-Eidechse (*L. fraasii*, Lehrs). Verlag nicht ermittelbar, *Zoologische Jahrbücher / Abteilung für Systematik*, 89, 407-478.
- Peterson, A. T. (2011). Ecological niche conservatism: A time-structured review of evidence. *Journal of Biogeography*, 38(5), 817–827.
- Peterson, A. T., Papeş, M., & Soberón, J. (2008). Rethinking receiver operating characteristic analysis applications in ecological niche modeling. *Ecological Modelling*, 213(1), 63–72.
- Petrosyan, V., Osipov, F., Bobrov, V., Dergunova, N., Nazarenko, E., Omelchenko, A., Danielyan, F., & Arakelyan, M. (2019). Analysis of geographical distribution of the parthenogenetic rock lizard *Darevskia armeniaca* and its parental species (*D. mixta*, *D. valentini*) based on ecological modelling. *Salamandra*, 55(3), 173–190.
- Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, 190(3–4), 231–259.
- Phillips, S. J., Dudík, M., & Schapire, R. E. (2017). Maxent software for modeling species niches and distributions (Version 3.4. 1). *Biodiversity Informatics*.
- Provan, J., & Bennett, K. D. (2008). Phylogeographic insights into cryptic glacial refugia. *Trends in Ecology & Evolution*, 23(10), 564–571.
- R. Core Team. (2018). R: A language and environment for statistical computing, R foundation for statistical computing. Vienna: R Core Team; 2018.
- Rivera, J. A., Rich, H. N., Michelle Lawing, A., Rosenberg, M. S., & Martins, E. P. (2021). Occurrence data uncover patterns of allopatric divergence and interspecies interactions in the evolutionary history of *Sceloporus* lizards. *Ecology and Evolution*, 11(6), 2796–2813.
- Rokas, A., Atkinson, R. J., Webster, L., Csóka, G., & Stone, G. N. (2003). Out of Anatolia: Longitudinal gradients in genetic diversity support an eastern origin for a circum-

- Mediterranean oak gallwasp *Andricus quercustozae*. *Molecular Ecology*, 12(8), 2153–2174.
- Sarıkaya, M. A., Zreda, M., Çiner, A., & Zweck, C. (2008). Cold and wet Last Glacial Maximum on Mount Sandıras, SW Turkey, inferred from cosmogenic dating and glacier modeling. *Quaternary Science Reviews*, 27(7–8), 769–780.
- Schmitt, T. (2009). Biogeographical and evolutionary importance of the European high mountain systems. *Frontiers in Zoology*, 6(1), 1–10.
- Schönswetter, P., Stehlik, I., Holderegger, R., & Tribsch, A. (2005). Molecular evidence for glacial refugia of mountain plants in the European Alps. *Molecular Ecology*, 14(11), 3547–3555.
- Şekercioğlu, Ç. H., Anderson, S., Akçay, E., Bilgin, R., Can, Ö. E., Semiz, G., Tavşanoğlu, Ç., Yokeş, M. B., Soyumert, A., & Ipekdağ, K. (2011). Turkey's globally important biodiversity in crisis. *Biological Conservation*, 144(12), 2752–2769.
- Sindaco, R., Venchi, A., Carpaneto, G. M., & Bologna, M. A. (2000). The reptiles of Anatolia: A checklist and zoogeographical analysis. *Biogeographia—The Journal of Integrative Biogeography*, 21(1), 441–554.
- Ülker, E. D., Tavşanoğlu, Ç., & Perktas, U. (2018). Ecological niche modelling of pedunculate oak (*Quercus robur*) supports the 'expansion–contraction' model of Pleistocene biogeography. *Biological Journal of the Linnean Society*, 123(2), 338–347.
- Unal, Y., Kindap, T., & Karaca, M. (2003). Redefining the climate zones of Turkey using cluster analysis. *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 23(9), 1045–1055.
- Varga, Z. (2009). Extra-Mediterranean refugia, post-glacial vegetation history and area dynamics in Eastern Central Europe. In *Relict species* (pp. 57–87). Springer.
- Waltari, E., Hijmans, R. J., Peterson, A. T., Nyári, A. S., Perkins, S. L., & Guralnick, R. P. (2007). Locating Pleistocene refugia: Comparing phylogeographic and ecological niche model predictions. *PLoS One*, 2(7), e563.
- Watanabe, M., Suzuki, T., O'ishi, R., Komuro, Y., Watanabe, S., Emori, S., Takemura, T., Chikira, M., Ogura, T., & Sekiguchi, M. (2010). Improved climate simulation by MIROC5: Mean states, variability, and climate sensitivity. *Journal of Climate*, 23(23), 6312–6335.
- [www.worldclim.org/](http://www.worldclim.org/). (n.d.). WorldClim - Global Climate Data | Free Climate Data for Ecological Modeling and GIS. Retrieved June 18, 2019, from <https://www.worldclim.org/>
- Zeng, Z.-G., Bi, J.-H., Li, S.-R., Chen, S.-Y., Pike, D. A., Gao, Y., & Du, W.-G. (2014). Effects of habitat alteration on lizard community and food web structure in a desert steppe ecosystem. *Biological Conservation*, 179, 86–92.